

Contents Volume 16, 1990

Editorial	v
A new planktic foraminifer transfer function for estimating Pliocene-Holocene paleoceanographic conditions in the North Atlantic	
H.J. Dowsett and R.Z. Poore (Reston, VA, U.S.A.)	1
Ecology and paleoecology of triserial planktic foraminifera	
D. Kroon and A.J. Nederbragt (Amsterdam, The Netherlands)	25
Coccoliths in sediment traps from the Norwegian Sea	
C. Samtleben and T. Bickert (Kiel, F.R.G.)	39
Distribution of coccoliths in surface sediments in the Norwegian-Greenland sea	
L.K. Eide (Bergen, Norway)	65
Response of Northern Indian Ocean deep-sea benthic foraminifera to global climates during Pliocene-Pleistocene	
A.K. Gupta (Kharagpur, India) and M.S. Srinivasan (Varanasi, India)	77
Effects of gametogenesis on test structure and dissolution of some spinose planktonic foraminifera and implications for test preservation	
D.A. Caron (Woods Hole, MA, U.S.A.), O.R. Anderson (Palisades, NY, U.S.A.), J.L. Lindsey (Fort Pierce, FL, U.S.A.), W.W. Faber Jr. (New York, NY, U.S.A.) and E.L. Lim (Woods Hole, MA, U.S.A.)	93
A quantitative micropaleontologic method for shallow marine paleoclimatology: Application to Pliocene deposits of the western North Atlantic	
T.M. Cronin and H.J. Dowsett (Reston, VA, U.S.A.)	117
"Squatter" behaviour in soft-shelled foraminifera	
L. Moodley (Den Burg, The Netherlands)	149
Experimental and observational studies of radiolarian physiological ecology: 4. Factors determining the distribution and survival of <i>Didymocyrtis tetrahalamus</i> tetrathalamus with implications for paleoecological interpretations	
O.R. Anderson, M. Bryan and P. Bennett (Palisades, NY, U.S.A.)	155
Recent obliquipithonelloid calcareous cysts of <i>Scrippsiella patagonica</i> sp. nov. (Peridiniaceae, Dinophyceae) from plankton of the Golfo San Jorge (Patagonia, Argentina)	
R. Akselman (Mar del Plata, Argentina) and H. Keupp (Berlin, G.D.R.)	169
Main features of foraminiferal distribution reflecting estuarine hydrography in Oslo Fjord	
E. Alve and J. Nagy (Oslo, Norway)	181
Comparison of fine fraction with monospecific foraminiferal stable isotopic stratigraphies from pelagic carbonates across the last glacial termination	
C.K. Paull (Chapel Hill, NC, U.S.A.) and H. Thierstein (Zürich, Switzerland)	207
Progenesis in Late Eocene populations of <i>Subbotina linaperta</i> (Foraminifera) from the western Atlantic	
N. MacLeod (Ann Arbor, MI, U.S.A.), G. Keller (Princeton, NJ, U.S.A.) and J.A. Kitchell (Ann Arbor, MI, U.S.A.)	219
Benthic foraminiferal assemblages from the eastern Weddell Sea between 68 and 73°S: Distribution, ecology and fossilization potential	
A. Mackensen, H. Grobe, G. Kuhn and D.K. Fütterer (Bremerhaven, F.R.G.)	241
Seasonal occurrence of silicoflagellate morphologies in different environments of the eastern Pacific Ocean	
C. Sancetta (Palisades, NY, U.S.A.)	285
Distribution and malformation of extant calcareous nannoplankton in the Indonesian Seas	
A. Kleijne (Amsterdam, The Netherlands)	293

